



**LAWN SOLUTIONS AUSTRALIA**

ACELEPRYN® is a registered trademark of a Syngenta Group Company

APVMA Approval No: 91933/ 133988

D.O.M. Batch



## READ SAFETY DIRECTIONS

**LAWN SOLUTIONS**

# Grub Guard Ultimate Turf Insecticide

**ACTIVE CONSTITUENT: 2G/KG CHLORANTRANILIPROLE**

For the control of major turf pests including Lawn Army Worm and Lawn Curl Grubs. Suitable for all lawn types



LAWN GRUBS



ARMY WORM



BLACK BEETLE

**POWERED BY ACELEPRYN®**

TREATS UP TO  
**4KG NET 500M<sup>2</sup>**



### HOW TO USE

LAWN SOLUTIONS Grub Guard Ultimate Turf Insecticide delivers excellent preventative control against a large range of common lawn pests. Optimal results are achieved when applied early in the season at the appearance of overwintering adult pests. The long residual performance of LAWN SOLUTIONS Grub Guard Ultimate Turf Insecticide will provide up to 6 months protection at the higher application rates. Curative control of pest outbreaks later in the season can also be achieved using the higher recommended application rates.

### Restrictions

DO NOT apply when the soil is saturated with water as adequate distribution of the product into the soil will not be achieved.

SITUATION	PEST	RATE	COMMENTS
All Lawns and Turfgrass	Lawn Curl Grub (Beetle larvae including: African Black Beetle larvae Argentinean Scarab larvae)	75 – 150 g per 10 m <sup>2</sup> or 750 g – 1.5 kg per 100 m <sup>2</sup>	African Black Beetle larvae Apply before or at peak egg hatch for maximum control (typically mid-September).  Argentinean Scarab larvae Apply before or at peak egg hatch for maximum control (typically mid-December).  Apply the higher rate for: • up to 6 months residual protection. • cases of high pest pressure • late in season when mature grubs are present for African Black Beetle (typically mid-December onwards)  Lightly water in immediately after application
	Argentine Stem Weevil larvae Billbug larvae		Apply mid-September to prevent damage and population build up. Early application is essential to prevent lawn damage due to larvae feeding.  Apply the high rate for: • up to 6 months residual protection in cases of high pest pressure. • late season when mature larvae are present, applications mid-December onwards  Lightly water in immediately after application
Caterpillars including Lawn Armyworm, Black Cutworm, and Sod Webworm		75 g per 10 m <sup>2</sup> or 750 g per 100 m <sup>2</sup>	Apply as a curative application at first sign of caterpillars or when damage is observed.  To ensure optimum control, delay watering or mowing for 24 hours after application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED**

### HOW TO APPLY

LAWN SOLUTIONS Grub Guard Ultimate Turf Insecticide is a ready to use spreadable insecticide granule. Measure the lawn area before application to determine size then weight amount of granules to apply over the area. Evenly spread granules over the lawn either by spreader or hand application.

**Spreader application:** The use of a garden spreader or professional spreader that can be calibrated for even output is recommended for optimal product performance. Use a drop type, rotary type or hand held spreader.

1. Weigh out the amount required to treat the measured area.
2. Fill spreader with granules.
3. Evenly apply granules over the lawn area.
4. Application may be made in one pass or multiple passes in different directions. When applying in multiple directions, separate the weighed out granules by the number of directions to be applied before treating each direction.

**Hand Application:** Hand application may be used to evenly broadcast granules over the lawn area. It is recommended to use gloves when applying by hand.

1. Weigh out the amount required to treat the measured area.
2. Fill container to hand broadcast granules from.
3. Evenly spread by hand, moving at a steady walking speed.
4. When hand applying it is recommended to apply with multiple passes in different directions to ensure even application. Separate the weighed out granules by the number of directions to be applied before treating each direction.

### After application wash in/irrigation

Lightly water in immediately after application for optimal control of beetle larvae (Black beetle, Argentine scarab, Argentine stem weevil and

Billbug). Delay watering and mowing for 24 hours after application when treating for caterpillars.

### Spreader Equipment Cleanout

Thoroughly clean spreader after application and other application equipment with clean water.

### Compatibility

As formulations of other manufacturer's granular products are beyond the control of Lawn Solutions, blends with other products are not recommended as products of differing size granules may result in uneven applications. For further product information visit [www.lawnsolutionsaustralia.com.au](http://www.lawnsolutionsaustralia.com.au)

### CAUTION

Toxic to aquatic life. DO NOT allow chemical containers or granules to get into drains, sewers, streams or ponds. DO NOT feed grass clippings from treated areas to poultry or other animals.

### STORAGE AND DISPOSAL

**KEEP OUT OF REACH OF CHILDREN**  
Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

### SAFETY DIRECTIONS

**May irritate the eyes, avoid contact with eyes. Wash hands after use.**

### FIRST AID

**First aid is not generally required. If in doubt, contact a doctor or Poisons Information Centre. Phone 131 126.**

**FOR EMERGENCIES ONLY CALL: 1800 033 111 (24 hrs)**

This product must be used strictly as directed. Not to be used for any use or in any manner contrary to the label.